

Casorati Determinant Solution for the Relativistic Toda Lattice Equation

Yasuhiro OHTA^{*}, Kenji KAJIWARA^a, Junta MATSUKIDAIRA^b
and Junkichi SATSUMA^c

*Research Institute for Mathematical Sciences,
Kyoto University, Kyoto 606, Japan*

^a*Department of Applied Physics, Faculty of Engineering,
University of Tokyo, Tokyo 113, Japan*

^b*Department of Applied Mathematics and Informatics,
Ryukoku University, Seta, Ohtsu 520-21, Japan*

^c*Department of Mathematical Sciences, University of Tokyo,
Meguro-ku, Tokyo 153, Japan*

The relativistic Toda lattice equation is decomposed into three Toda systems, the Toda lattice itself, Bäcklund transformation of Toda lattice and discrete time Toda lattice. It is shown that the solutions of the equation are given in terms of the Casorati determinant. By using the Casoratian technique, the bilinear equations of Toda systems are reduced to the Laplace expansion form for determinants. The N -soliton solution is explicitly constructed in the form of the Casorati determinant.

^{*} On leave from Department of Applied Mathematics, Faculty of Engineering, Hiroshima University.

I. INTRODUCTION

The relativistic Toda lattice (RT) equation,

$$\ddot{q}_n = (1 + \frac{1}{c}\dot{q}_{n-1})(1 + \frac{1}{c}\dot{q}_n) \frac{\exp(q_{n-1} - q_n)}{1 + \frac{1}{c^2}\exp(q_{n-1} - q_n)} - (1 + \frac{1}{c}\dot{q}_n)(1 + \frac{1}{c}\dot{q}_{n+1}) \frac{\exp(q_n - q_{n+1})}{1 + \frac{1}{c^2}\exp(q_n - q_{n+1})}, \quad (1.1)$$

where q_n is the coordinates of n -th lattice point, and $\dot{}$ means the differentiation with respect to time t and c is the light speed, was introduced and studied by Ruijsenaars.¹ This equation is derived by requiring the Poincaré invariance to the well-known Toda lattice (TL) equation. The action-angle transformation and complete integrability were also investigated for the RT equation. Bruschi and Ragnisco² constructed the recursion operator and Bäcklund transformation for eq. (1.1). The relation between the discrete time Toda lattice (DTL) and RT was found by Suris³ in the context of the Hamiltonian dynamics. We will show later that the DTL equation actually takes a part of the RT equation. In the limit of $c \rightarrow \infty$, the RT eq. (1.1) reduces to the ordinary TL equation,

$$\ddot{q}_n = \exp(q_{n-1} - q_n) - \exp(q_n - q_{n+1}), \quad (1.2)$$

which is one of the most typical equations in soliton theory.

The bilinear formalism introduced by Hirota⁴ is a key to understand the algebraic structure of soliton equations. By variable transformations, soliton equations are transformed into the bilinear form. The solutions for the bilinear equations are given in terms of certain determinants or Pfaffians. We analyze the RT equation from the view point of the bilinear theory. The Wronski determinant representation of solutions was first given for the KdV equation.⁵ Freeman and Nimmo⁶ have developed the Laplace expansion technique and given the Wronskian solutions for various soliton equations. Hirota, Ito and Kako⁷ showed that for the discrete analogues of soliton equations, solutions for their bilinear form are given by the Casorati determinant instead of Wronskian.

M. and Y. Sato^{8,9} found that the soliton equations are reduced to the Plücker relation with respect to the τ function which is expressed by the infinite dimensional Wronski determinant. Date, Jimbo, Kashiwara and Miwa^{10,11} developed the theory of transformation group for solutions of soliton equations. Discrete soliton equations were also treated in their context.^{11,12} Ueno and Takasaki¹³ extended Sato's idea to the Toda lattice hierarchy.

Recently Hietarinta and one of the authors (J.S.) found that the RT equation is transformed into a trilinear form through a suitable dependent variable transformation.¹⁴ It was also shown that solutions for the trilinear equation are given by the determinant with Wronskian structure in two directions, rowwise and columnwise. The trilinear equation can be regarded as an extension of the bilinear form. Therefore it may be expected that the trilinear form has abundant algebraic structure.¹⁵ However the trilinear equation for RT is in the case that the coupling constant is negative, and hence the situation is not considered to be physical. To overcome this difficulty, we have investigated the solutions of eq. (1.1) and it has become clear that this equation holds various algebraic structures simultaneously.

In this paper, we propose another type of determinant solution to the RT eq. (1.1), the Casorati determinant. This determinant represents the N -soliton solution for RT, so fits for the physical situation. The solution converges to the Wronskian solution for the one-dimensional TL equation in the limit of $c \rightarrow \infty$. The RT equation is transformed into three bilinear equations, the TL equation, the Bäcklund transformation (BT) for TL and the DTL equation. We show how the RT equation is composed of these three Toda systems. It is proved by using the Laplace expansion and the reduction technique that the Casorati determinant satisfies the bilinear equations. For discrete soliton equations, the difference formula is simply obtained by using the Casoratian technique.¹⁶ Through the differential formula and difference formula for the Casorati determinant, the bilinear equations of RT are reduced to the “algebraic” identities. In **II**, the bilinearization of the RT equation is carried out by introducing the auxiliary dependent variables. In **III**, we show the Casorati determinant solution for the bilinear equations which represents the N -soliton solution. A constructive proof that the determinant satisfies the RT equation is given in **IV**, and **V**. At first in **IV**, the solutions for the two-dimensional Toda lattice (2DTL) equation, BT of TL and discrete time two-dimensional Toda lattice (D2DTL) equation are derived in the form of the Wronski and Casorati determinants. Next in **V**, we explain the reduction technique which reduces the 2DTL and D2DTL equations into the TL and DTL equations, respectively. In **VI**, concluding remarks are given.

II. BILINEAR FORM FOR THE RT EQUATION

The RT eq. (1.1) is transformed into the three bilinear equations,

$$D_x^2 f_n \cdot f_n = 2(\bar{g}_n g_n - f_n^2), \quad (2.1a)$$

$$(aD_x - 1)f_n \cdot f_{n-1} + \bar{g}_{n-1}g_n = 0, \quad (2.1b)$$

$$\bar{g}_{n-1}g_{n+1} - f_n^2 = a^2(f_{n+1}f_{n-1} - f_n^2), \quad (2.1c)$$

through the variable transformations,

$$q_n = \log \frac{f_{n-1}}{f_n}, \quad (2.2a)$$

$$t = \frac{\sqrt{1+c^2}}{c}x, \quad (2.2b)$$

$$c = \frac{\sqrt{1-a^2}}{a}. \quad (2.2c)$$

In eqs. (2.1), g and $\bar{g}$ are the auxiliary variables and D is the bilinear differential operator defined by⁴

$$D_x^k f \cdot g = (\partial_x - \partial_y)^k f(x)g(y)|_{y=x}. \quad (2.3)$$

As will be shown later, eqs. (2.1a), (2.1b) and (2.1c) are nothing but the bilinear forms of TL equation, BT of TL and DTL equation, respectively.

Let us show that the RT equation actually follows from the bilinear eqs. (2.1). Dividing eqs. (2.1a), (2.1b) and (2.1c) by $2f_n^2$, $f_{n-1}f_n$ and $f_{n-1}f_{n+1}$ respectively, we get

$$\frac{\bar{g}_n g_n}{f_n f_n} = 1 + (\log f_n)_{xx}, \quad (2.4a)$$

$$\frac{\bar{g}_{n-1} g_n}{f_{n-1} f_n} = 1 + a(\log \frac{f_{n-1}}{f_n})_x, \quad (2.4b)$$

$$\frac{\bar{g}_{n-1} g_{n+1}}{f_{n-1} f_{n+1}} = a^2 + (1-a^2) \frac{f_n f_n}{f_{n-1} f_{n+1}}, \quad (2.4c)$$

where the subscript x indicates differentiation. We have an identity

$$\frac{\bar{g}_{n-1} g_{n+1}}{f_{n-1} f_{n+1}} \frac{\bar{g}_n g_n}{f_n f_n} = \frac{\bar{g}_{n-1} g_n}{f_{n-1} f_n} \frac{\bar{g}_n g_{n+1}}{f_n f_{n+1}}, \quad (2.5)$$

which is rewritten by using eqs. (2.4) as

$$(a^2 + (1-a^2) \frac{f_n f_n}{f_{n-1} f_{n+1}}) (1 + (\log f_n)_{xx}) = (1 + a(\log \frac{f_{n-1}}{f_n})_x) (1 + a(\log \frac{f_n}{f_{n+1}})_x). \quad (2.6)$$

Noticing

$$\frac{1-a^2}{a^2} = c^2, \quad (2.7)$$

$$\partial_x = \frac{\sqrt{1+c^2}}{c} \partial_t = \frac{1}{ac} \partial_t, \quad (2.8)$$

we get from eq. (2.6)

$$a^2 + \frac{1}{c^2} (\log f_n)_{tt} = (1 + \frac{1}{c} \dot{q}_n) (1 + \frac{1}{c} \dot{q}_{n+1}) / (1 + c^2 \exp(q_{n+1} - q_n)). \quad (2.9)$$

Finally subtracting eq. (2.9) from eq. (2.9) with n replaced by $n-1$, we obtain

$$\begin{aligned} \frac{1}{c^2} \ddot{q}_n &= (1 + \frac{1}{c} \dot{q}_{n-1}) (1 + \frac{1}{c} \dot{q}_n) / (1 + c^2 \exp(q_n - q_{n-1})) \\ &\quad - (1 + \frac{1}{c} \dot{q}_n) (1 + \frac{1}{c} \dot{q}_{n+1}) / (1 + c^2 \exp(q_{n+1} - q_n)), \end{aligned} \quad (2.10)$$

which is the RT eq. (1.1) itself. Therefore if f_n , g_n and $\bar{g}_n$ satisfy eqs. (2.1), q_n defined by eq. (2.2a) gives the solution for RT equation.

III. CASORATI DETERMINANT SOLUTION FOR THE RT EQUATION

The Casorati determinant representation of soliton solution for the bilinear RT eq. (2.1) is given by

$$f_n = \begin{vmatrix} \varphi_1^{(n)} & \varphi_1^{(n+1)} & \cdots & \varphi_1^{(n+N-1)} \\ \varphi_2^{(n)} & \varphi_2^{(n+1)} & \cdots & \varphi_2^{(n+N-1)} \\ \vdots & \vdots & & \vdots \\ \varphi_N^{(n)} & \varphi_N^{(n+1)} & \cdots & \varphi_N^{(n+N-1)} \end{vmatrix}, \quad (3.1a)$$

$$g_n = \begin{vmatrix} \psi_1^{(n)} & \psi_1^{(n+1)} & \cdots & \psi_1^{(n+N-1)} \\ \psi_2^{(n)} & \psi_2^{(n+1)} & \cdots & \psi_2^{(n+N-1)} \\ \vdots & \vdots & & \vdots \\ \psi_N^{(n)} & \psi_N^{(n+1)} & \cdots & \psi_N^{(n+N-1)} \end{vmatrix}, \quad (3.1b)$$

$$\bar{g}_n = \begin{vmatrix} \bar{\psi}_1^{(n)} & \bar{\psi}_1^{(n+1)} & \cdots & \bar{\psi}_1^{(n+N-1)} \\ \bar{\psi}_2^{(n)} & \bar{\psi}_2^{(n+1)} & \cdots & \bar{\psi}_2^{(n+N-1)} \\ \vdots & \vdots & & \vdots \\ \bar{\psi}_N^{(n)} & \bar{\psi}_N^{(n+1)} & \cdots & \bar{\psi}_N^{(n+N-1)} \end{vmatrix}, \quad (3.1c)$$

and

$$\varphi_i^{(n)} = \left(\frac{p_i}{1 - p_i a} \right)^n e^{\eta_i} + \left(\frac{p_i^{-1}}{1 - p_i^{-1} a} \right)^n e^{\xi_i}, \quad (3.2a)$$

$$\psi_i^{(n)} = p_i^{-1} \left(\frac{p_i}{1 - p_i a} \right)^n e^{\eta_i} + p_i \left(\frac{p_i^{-1}}{1 - p_i^{-1} a} \right)^n e^{\xi_i}, \quad (3.2b)$$

$$\bar{\psi}_i^{(n)} = p_i \left(\frac{p_i}{1 - p_i a} \right)^n e^{\eta_i} + p_i^{-1} \left(\frac{p_i^{-1}}{1 - p_i^{-1} a} \right)^n e^{\xi_i}, \quad (3.2c)$$

$$\eta_i = p_i x + \eta_{i0}, \quad (3.3a)$$

$$\xi_i = p_i^{-1} x + \xi_{i0}, \quad (3.3b)$$

where N is the number of solitons, and p_i and $\eta_{i0} - \xi_{i0}$ are the wave number and phase parameter of i -th soliton respectively.

In the non-relativistic limit, c tends to infinity, a becomes 0 and

$$g_{n+1} = \bar{g}_{n-1} = f_n, \quad (3.4)$$

$$x = t. \quad (3.5)$$

In this case, eq. (2.1a) becomes the bilinear form of TL equation,

$$D_t^2 f_n \cdot f_n = 2(f_{n+1} f_{n-1} - f_n^2), \quad (3.6)$$

and eqs. (2.1b) and (2.1c) are trivial. Equation (3.6) is transformed into the TL eq. (1.2) through the variable transformation (2.2a). At $c \rightarrow \infty$, the determinant f_n in eq. (3.1a) gives the N -soliton solution to the TL equation in Wronskian form.

We will prove by using the Wronskian technique and Casoratian technique that the bilinear eqs. (2.1) are satisfied by f_n , g_n and $\bar{g}_n$ in eqs. (3.1). At first we give the determinant solution for the 2DTL, BT of TL and D2DTL equations in **IV**. Next in **V**, applying the reduction technique we recover the Casorati determinant solution in eqs. (3.1).

IV. THREE TODA SYSTEMS AND THEIR DETERMINANT SOLUTIONS

In this section we show that the solutions for the 2DTL equation, BT of TL and D2DTL equation are given in terms of the Wronskian, Casorati determinant and also Casorati determinant, respectively.

A. 2DTL

The bilinear form of 2DTL equation is written as

$$D_{x_1} D_{x_{-1}} \tau_n \cdot \tau_n = 2(\tau_n \tau_n - \tau_{n+1} \tau_{n-1}). \quad (4.1)$$

The Wronskian solution for this equation is given by^{7,17}

$$\tau_n = \begin{vmatrix} \varphi_1^{(n)} & \varphi_1^{(n+1)} & \cdots & \varphi_1^{(n+N-1)} \\ \varphi_2^{(n)} & \varphi_2^{(n+1)} & \cdots & \varphi_2^{(n+N-1)} \\ \vdots & \vdots & & \vdots \\ \varphi_N^{(n)} & \varphi_N^{(n+1)} & \cdots & \varphi_N^{(n+N-1)} \end{vmatrix}, \quad (4.2)$$

where $\varphi_i^{(n)}$'s are arbitrary functions satisfying the dispersion relations,

$$\partial_{x_\nu} \varphi_i^{(n)} = \varphi_i^{(n+\nu)}, \quad \nu = 1, -1. \quad (4.3)$$

After Freeman and Nimmo,⁶ we use the following notation for simplicity,

$$|n_1, n_2, \dots, n_N| = \begin{vmatrix} \varphi_1^{(n_1)} & \varphi_1^{(n_2)} & \cdots & \varphi_1^{(n_N)} \\ \varphi_2^{(n_1)} & \varphi_2^{(n_2)} & \cdots & \varphi_2^{(n_N)} \\ \vdots & \vdots & & \vdots \\ \varphi_N^{(n_1)} & \varphi_N^{(n_2)} & \cdots & \varphi_N^{(n_N)} \end{vmatrix}. \quad (4.4)$$

In this notation, τ_n in eq. (4.2) is rewritten as

$$\tau_n = |n, n+1, \dots, n+N-1|. \quad (4.5)$$

We show that the above Wronskian satisfies the bilinear eq. (4.1) by using the Wronskian technique. From eq. (4.3), the differentiation of the determinant $|n_1, n_2, \dots, n_N|$ is given by

$$\partial_{x_\nu} |n_1, n_2, \dots, n_N| = \sum_{j=1}^N |n_1, n_2, \dots, n_j + \nu, \dots, n_N|, \quad \nu = 1, -1. \quad (4.6)$$

Thus we get the following differential formula for τ_n ,

$$\partial_{x_1} \tau_n = |n, n+1, \dots, n+N-2, n+N|, \quad (4.7a)$$

$$\partial_{x_{-1}} \tau_n = |n-1, n+1, \dots, n+N-2, n+N-1|, \quad (4.7b)$$

$$\begin{aligned} \partial_{x_1} \partial_{x_{-1}} \tau_n &= |n-1, n+1, \dots, n+N-2, n+N| \\ &\quad + |n, n+1, \dots, n+N-2, n+N-1|. \end{aligned} \quad (4.7c)$$

Now we consider an identity for $2N \times 2N$ determinant,

$$\begin{vmatrix} n-1 & n+1 & \cdots & n+N-2 & n+N & n & \cdots & n+N-1 \\ \hline n-1 & & \cdots & \emptyset & n & n+1 & \cdots & n+N-1 \end{vmatrix} = 0, \quad (4.8)$$

where a number denotes a column vector of height N and $\emptyset$ means an empty block. Applying the Laplace expansion for determinant to the left-hand side of above equation, we obtain the algebraic bilinear identity for determinants,

$$\begin{aligned} & |n-1, n+1, \dots, n+N-2, n+N| |n, n+1, \dots, n+N-2, n+N-1| \\ & - |n, n+1, \dots, n+N-2, n+N| |n-1, n+1, \dots, n+N-2, n+N-1| \\ & + |n+1, \dots, n+N-2, n+N-1, n+N| |n-1, n, n+1, \dots, n+N-2| = 0. \end{aligned} \quad (4.9)$$

By using eqs. (4.7), this equation is rewritten into the differential bilinear equation for τ_n ,

$$(\partial_{x_1} \partial_{x_{-1}} \tau_n - \tau_n) \tau_n - \partial_{x_1} \tau_n \partial_{x_{-1}} \tau_n + \tau_{n+1} \tau_{n-1} = 0, \quad (4.10)$$

which recovers eq. (4.1). Hence we have proved that the Wronskian τ_n gives the solution for 2DTL equation.

B. BT of TL

We have the bilinear form for BT of TL equation,¹⁸

$$(aD_x - 1)\tau'_n \cdot \tau_{n-1} + \tau_n \tau'_{n-1} = 0, \quad (4.11)$$

where a is a constant and x corresponds to x_1 in eq. (4.1), which transforms a TL τ_n into another TL τ'_n . Applying the BT recursively, we get an infinite sequence of TL's, $\tau_n, \tau'_n, \tau''_n, \dots$. We denote the k -th TL as $\tau_n(k)$. Then the BT of TL is written as

$$(aD_x - 1)\tau_n(k+1) \cdot \tau_{n-1}(k) + \tau_n(k)\tau_{n-1}(k+1) = 0. \quad (4.12)$$

The Casorati determinant solution for eq. (4.12) is given by

$$\tau_n(k) = \begin{vmatrix} \varphi_1^{(n)}(k) & \varphi_1^{(n+1)}(k) & \dots & \varphi_1^{(n+N-1)}(k) \\ \varphi_2^{(n)}(k) & \varphi_2^{(n+1)}(k) & \dots & \varphi_2^{(n+N-1)}(k) \\ \vdots & \vdots & \dots & \vdots \\ \varphi_N^{(n)}(k) & \varphi_N^{(n+1)}(k) & \dots & \varphi_N^{(n+N-1)}(k) \end{vmatrix}, \quad (4.13)$$

where $\varphi_i^{(n)}(k)$'s are arbitrary functions satisfying the dispersion relations,

$$\partial_x \varphi_i^{(n)}(k) = \varphi_i^{(n+1)}(k), \quad (4.14a)$$

$$\Delta_k \varphi_i^{(n)}(k) = \varphi_i^{(n+1)}(k), \quad (4.14b)$$

where Δ_k is the backward difference operator with the difference interval a defined by

$$\Delta_k f(k) = \frac{f(k) - f(k-1)}{a}. \quad (4.15)$$

For simplicity we introduce a convenient notation,

$$|n_{1k_1}, n_{2k_2}, \dots, n_{Nk_N}| = \begin{vmatrix} \varphi_1^{(n_1)}(k_1) & \varphi_1^{(n_2)}(k_2) & \cdots & \varphi_1^{(n_N)}(k_N) \\ \varphi_2^{(n_1)}(k_1) & \varphi_2^{(n_2)}(k_2) & \cdots & \varphi_2^{(n_N)}(k_N) \\ \vdots & \vdots & & \vdots \\ \varphi_N^{(n_1)}(k_1) & \varphi_N^{(n_2)}(k_2) & \cdots & \varphi_N^{(n_N)}(k_N) \end{vmatrix}. \quad (4.16)$$

In this notation, the solution for BT of TL, $\tau_n(k)$ in eq. (4.13), is rewritten as

$$\tau_n(k) = |n_k, n+1_k, \dots, n+N-1_k|, \quad (4.17)$$

or suppressing the index k ,

$$\tau_n(k) = |n, n+1, \dots, n+N-1|. \quad (4.18)$$

We show that the above $\tau_n(k)$ actually satisfies the bilinear eq. (4.12). At first we investigate the difference formula for τ . From eq. (4.14b) we have

$$a\varphi_i^{(n+1)}(k+1) = \varphi_i^{(n)}(k+1) - \varphi_i^{(n)}(k). \quad (4.19)$$

Noticing this relation, we get

$$\begin{aligned} \tau_n(k+1) &= |n_{k+1}, n+1_{k+1}, n+2_{k+1}, \dots, n+N-1_{k+1}| \\ &= |n_k, n+1_{k+1}, n+2_{k+1}, \dots, n+N-1_{k+1}| \end{aligned}$$

where we have subtracted the 2nd column multiplied by a from the 1st column,

$$= |n_k, n+1_k, n+2_{k+1}, \dots, n+N-1_{k+1}|$$

where we have subtracted the 3rd column multiplied by a from the 2nd column,

...

$$= |n_k, n+1_k, \dots, n+N-2_k, n+N-1_{k+1}|,$$

that is,

$$\tau_n(k+1) = |n_k, n+1_k, \dots, n+N-2_k, n+N-1_{k+1}|, \quad (4.20)$$

or suppressing the index k ,

$$\tau_n(k+1) = |n, n+1, \dots, n+N-2, n+N-1_{k+1}|. \quad (4.21)$$

Moreover in eq. (4.20), multiplying the N -th column by a and adding the $(N-1)$ -th column to the N -th column, we get

$$a\tau_n(k+1) = |n_k, n+1_k, \dots, n+N-2_k, n+N-2_{k+1}|. \quad (4.22)$$

The important point is that the effect of difference appears only one column as shown in eqs. (4.20) and (4.22). Consequently the Laplace expansion becomes applicable. This technique for condensing the shift of index into only one column may be called a Casoratian technique.

Differentiating eq. (4.22) with x and using eq. (4.14a) we obtain

$$\begin{aligned} a\partial_x \tau_n(k+1) &= |n_k, n+1_k, \dots, n+N-3_k, n+N-1_k, n+N-2_{k+1}| \\ &\quad + |n_k, n+1_k, \dots, n+N-3_k, n+N-2_k, n+N-1_{k+1}| \\ &= |n_k, n+1_k, \dots, n+N-3_k, n+N-1_k, n+N-2_{k+1}| \\ &\quad + \tau_n(k+1), \end{aligned} \quad (4.23)$$

where we have used eq. (4.20) in the last line. We have also

$$\partial_x \tau_{n-1}(k) = |n-1, n, \dots, n+N-3, n+N-1|. \quad (4.24)$$

Let us introduce an identity for $2N \times 2N$ determinant,

$$\left| \begin{array}{ccc|ccc|cccc|c} n_k & \cdots & n+N-3_k & n+N-1_k & n+N-2_{k+1} & & & \emptyset & & n+N-2_k \\ \hline & & \emptyset & & & & & & & \end{array} \right| = 0. \quad (4.25)$$

Applying the Laplace expansion to the left-hand side, we obtain the algebraic bilinear identity for determinants,

$$\begin{aligned} &|n_k, \dots, n+N-3_k, n+N-1_k, n+N-2_{k+1}| |n-1_k, n_k, \dots, n+N-3_k, n+N-2_k| \\ &- |n_k, \dots, n+N-3_k, n+N-2_k, n+N-2_{k+1}| |n-1_k, n_k, \dots, n+N-3_k, n+N-1_k| \\ &- |n_k, \dots, n+N-3_k, n+N-1_k, n+N-2_k| |n-1_k, n_k, \dots, n+N-3_k, n+N-2_{k+1}| = 0, \end{aligned} \quad (4.26)$$

which is rewritten by using eqs. (4.20), (4.22), (4.23) and (4.24), into the differential bilinear equation,

$$(a\partial_x\tau_n(k+1) - \tau_n(k+1))\tau_{n-1}(k) - a\tau_n(k+1)\partial_x\tau_{n-1}(k) + \tau_n(k)\tau_{n-1}(k+1) = 0. \quad (4.27)$$

This equation is equal to eq. (4.12). Hence we have proved that the Casorati determinant $\tau_n(k)$ in eq. (4.13) gives the solution for the BT of TL equation.

C. D2DTL

Using the Casoratian technique, we show that the D2DTL equation is satisfied by the Casorati determinant. The bilinear form of D2DTL equation is given by^{7,19}

$$\begin{aligned} & \tau_n(k+1, l+1)\tau_n(k, l) - \tau_n(k+1, l)\tau_n(k, l+1) \\ & = ab(\tau_n(k+1, l+1)\tau_n(k, l) - \tau_{n+1}(k+1, l)\tau_{n-1}(k, l+1)), \end{aligned} \quad (4.28)$$

where τ_n depends on two discrete independent variables k and l , and a and b are the difference intervals for k and l respectively.

The Casorati determinant solution for eq. (4.28) is given as

$$\tau_n(k, l) = \begin{vmatrix} \varphi_1^{(n)}(k, l) & \varphi_1^{(n+1)}(k, l) & \cdots & \varphi_1^{(n+N-1)}(k, l) \\ \varphi_2^{(n)}(k, l) & \varphi_2^{(n+1)}(k, l) & \cdots & \varphi_2^{(n+N-1)}(k, l) \\ \vdots & \vdots & & \vdots \\ \varphi_N^{(n)}(k, l) & \varphi_N^{(n+1)}(k, l) & \cdots & \varphi_N^{(n+N-1)}(k, l) \end{vmatrix}, \quad (4.29)$$

where $\varphi_i^{(n)}$'s are arbitrary functions of k and l which satisfy the dispersion relations,

$$\Delta_k \varphi_i^{(n)}(k, l) = \varphi_i^{(n+1)}(k, l), \quad (4.30a)$$

$$\Delta_l \varphi_i^{(n)}(k, l) = \varphi_i^{(n-1)}(k, l). \quad (4.30b)$$

Here Δ_k and Δ_l are the backward difference operators defined by

$$\Delta_k f(k, l) = \frac{f(k, l) - f(k-1, l)}{a}, \quad (4.31a)$$

$$\Delta_l f(k, l) = \frac{f(k, l) - f(k, l-1)}{b}. \quad (4.31b)$$

We use the following simple notation similar to that in eq. (4.16),

$$|n_{1 \underset{l_1}{k_1}}, n_{2 \underset{l_2}{k_2}}, \dots, n_{N \underset{l_N}{k_N}}| = \begin{vmatrix} \varphi_1^{(n_1)}(k_1, l_1) & \varphi_1^{(n_2)}(k_2, l_2) & \cdots & \varphi_1^{(n_N)}(k_N, l_N) \\ \varphi_2^{(n_1)}(k_1, l_1) & \varphi_2^{(n_2)}(k_2, l_2) & \cdots & \varphi_2^{(n_N)}(k_N, l_N) \\ \vdots & \vdots & & \vdots \\ \varphi_N^{(n_1)}(k_1, l_1) & \varphi_N^{(n_2)}(k_2, l_2) & \cdots & \varphi_N^{(n_N)}(k_N, l_N) \end{vmatrix}. \quad (4.32)$$

In this notation, the solution to D2DTL equation in eq. (4.29) is

$$\tau_n(k, l) = |n_{\underset{l}{k}}, n + 1_{\underset{l}{k}}, \dots, n + N - 1_{\underset{l}{k}}|. \quad (4.33)$$

Neglecting the index k and l , we write

$$\tau_n(k, l) = |n, n + 1, \dots, n + N - 1|. \quad (4.34)$$

Now let us examine the difference formula for the above τ_n . Note that the dispersion relation for k in eq. (4.31a) is equal to the one for the solution to BT of TL in **B**. Hence in the same manner as shown in eqs. (4.20) and (4.22), we obtain

$$\tau_n(k + 1, l) = |n_{\underset{l}{k}}, n + 1_{\underset{l}{k}}, \dots, n + N - 2_{\underset{l}{k}}, n + N - 1_{\underset{l}{k+1}}|, \quad (4.35)$$

$$a\tau_n(k + 1, l) = |n_{\underset{l}{k}}, n + 1_{\underset{l}{k}}, \dots, n + N - 2_{\underset{l}{k}}, n + N - 2_{\underset{l}{k+1}}|. \quad (4.36)$$

Similarly we get

$$\tau_n(k, l + 1) = |n_{\underset{l+1}{k}}, n + 1_{\underset{l}{k}}, \dots, n + N - 2_{\underset{l}{k}}, n + N - 1_{\underset{l}{k}}|, \quad (4.37)$$

$$b\tau_n(k, l + 1) = |n + 1_{\underset{l+1}{k}}, n + 1_{\underset{l}{k}}, \dots, n + N - 2_{\underset{l}{k}}, n + N - 1_{\underset{l}{k}}|, \quad (4.38)$$

because the role of l and b is parallel to that of k and a except that the ordering of index n is reversed. In short, we write

$$\tau_n(k + 1, l) = |n, n + 1, \dots, n + N - 2, n + N - 1_{k+1}|, \quad (4.39)$$

$$a\tau_n(k + 1, l) = |n, n + 1, \dots, n + N - 2, n + N - 2_{k+1}|, \quad (4.40)$$

$$\tau_n(k, l + 1) = |n_{l+1}, n + 1, \dots, n + N - 2, n + N - 1|, \quad (4.41)$$

$$b\tau_n(k, l + 1) = |n + 1_{l+1}, n + 1, \dots, n + N - 2, n + N - 1|. \quad (4.42)$$

We give the Casoratian technique for two variable shifted case. From eq. (4.41), $\tau_n(k+1, l+1)$ is given by

$$\tau_n(k+1, l+1) = |n_{\substack{k+1 \\ l+1}}, n+1_{k+1}, \dots, n+N-2_{k+1}, n+N-1_{k+1}|. \quad (4.43)$$

In the following we show that the shifts of index k is condensed into only the most right column of the determinant. From eqs. (4.30), $\varphi_i^{(n)}$ satisfies

$$\Delta_k \Delta_l \varphi_i^{(n)} = \varphi_i^{(n)}, \quad (4.44)$$

that is,

$$\varphi_i^{(n)}(k, l) - \varphi_i^{(n)}(k, l-1) - \varphi_i^{(n)}(k-1, l) + \varphi_i^{(n)}(k-1, l-1) = ab\varphi_i^{(n)}(k, l), \quad (4.45)$$

which is rewritten as

$$\begin{aligned} (1-ab)\varphi_i^{(n)}(k+1, l+1) &= \varphi_i^{(n)}(k+1, l) + \varphi_i^{(n)}(k, l+1) - \varphi_i^{(n)}(k, l) \\ &= \varphi_i^{(n)}(k, l+1) + a\varphi_i^{(n+1)}(k+1, l). \end{aligned} \quad (4.46)$$

Therefore multiplying the both hand sides of eq. (4.43) by $(1-ab)$ and rewriting the 1st column of the determinant by the use of eq. (4.46), we obtain

$$\begin{aligned} (1-ab)\tau_n(k+1, l+1) &= |n_{\substack{k \\ l+1}}, n+1_{k+1}, \dots, n+N-2_{k+1}, n+N-1_{k+1}| \\ &\quad + a|n+1_{\substack{k+1 \\ l}}, n+1_{k+1}, \dots, n+N-2_{k+1}, n+N-1_{k+1}|. \end{aligned} \quad (4.47)$$

The second term in r.h.s. vanishes because its 1st and 2nd columns are the same. So we get

$$(1-ab)\tau_n(k+1, l+1) = |n_{l+1}, n+1_{k+1}, \dots, n+N-2_{k+1}, n+N-1_{k+1}|. \quad (4.48)$$

In the above determinant, by subtracting the $(j+1)$ -th column multiplied by a from the j -th column for $j = 2, 3, \dots, N-1$, $\tau_n(k+1, l+1)$ is given by

$$(1-ab)\tau_n(k+1, l+1) = |n_{l+1}, n+1, \dots, n+N-2, n+N-1_{k+1}|. \quad (4.49)$$

As is shown, even if the two variables k and l are shifted, τ_n is also expressed in the determinant form whose columns are almost unchanged and only edges are varied. Thus we can use the Laplace expansion technique.

We consider an identity for $2N \times 2N$ determinant,

$$\begin{vmatrix} n_{l+1} & n+1 & \cdots & n+N-2 & n+N-1_{k+1} & n & \emptyset & n+N-1 \\ \hline n_{l+1} & \emptyset & & & & n & n+1 & \cdots & n+N-2 & n+N-1 \end{vmatrix} = 0. \quad (4.50)$$

By the Laplace expansion, we have

$$\begin{aligned} & |n_{l+1}, n+1, \dots, n+N-2, n+N-1_{k+1}| |n, n+1, \dots, n+N-2, n+N-1| \\ & - |n, n+1, \dots, n+N-2, n+N-1_{k+1}| |n_{l+1}, n+1, \dots, n+N-2, n+N-1| \\ & + |n+1, \dots, n+N-2, n+N-1, n+N-1_{k+1}| |n_{l+1}, n, n+1, \dots, n+N-2| = 0. \end{aligned} \quad (4.51)$$

Substituting eqs. (4.39)-(4.42) and (4.49) into eq. (4.51), we obtain

$$\begin{aligned} & (1-ab)\tau_n(k+1, l+1)\tau_n(k, l) - \tau_n(k+1, l)\tau_n(k, l+1) \\ & + a\tau_{n+1}(k+1, l)b\tau_{n-1}(k, l+1) = 0, \end{aligned} \quad (4.52)$$

which recovers eq. (4.28). This completes the proof that the Casorati determinant is indeed the solution for D2DTL equation.

V. REDUCTION TO THE RT EQUATION

Summarizing the arguments in **IV**, we see that the three bilinear equations,

$$D_x D_y \tau_n(k, l) \cdot \tau_n(k, l) = 2(\tau_n(k, l)\tau_n(k, l) - \tau_{n+1}(k, l)\tau_{n-1}(k, l)), \quad (5.1a)$$

$$(aD_x - 1)\tau_n(k+1, l) \cdot \tau_{n-1}(k, l) + \tau_n(k, l)\tau_{n-1}(k+1, l) = 0, \quad (5.1b)$$

$$\begin{aligned} & \tau_n(k+1, l+1)\tau_n(k, l) - \tau_n(k+1, l)\tau_n(k, l+1) \\ & = ab(\tau_n(k+1, l+1)\tau_n(k, l) - \tau_{n+1}(k+1, l)\tau_{n-1}(k, l+1)), \end{aligned} \quad (5.1c)$$

are simultaneously satisfied by the Casorati determinant,

$$\tau_n(k, l) = \begin{vmatrix} \varphi_1^{(n)}(k, l) & \varphi_1^{(n+1)}(k, l) & \cdots & \varphi_1^{(n+N-1)}(k, l) \\ \varphi_2^{(n)}(k, l) & \varphi_2^{(n+1)}(k, l) & \cdots & \varphi_2^{(n+N-1)}(k, l) \\ \vdots & \vdots & & \vdots \\ \varphi_N^{(n)}(k, l) & \varphi_N^{(n+1)}(k, l) & \cdots & \varphi_N^{(n+N-1)}(k, l) \end{vmatrix}, \quad (5.2)$$

supposed that $\varphi_i^{(n)}$'s obey the dispersion relations,

$$\partial_x \varphi_i^{(n)} = \varphi_i^{(n+1)}, \quad (5.3a)$$

$$\partial_y \varphi_i^{(n)} = \varphi_i^{(n-1)}, \quad (5.3b)$$

$$\Delta_k \varphi_i^{(n)} = \varphi_i^{(n+1)}, \quad (5.3c)$$

$$\Delta_l \varphi_i^{(n)} = \varphi_i^{(n-1)}, \quad (5.3d)$$

where Δ_k and Δ_l are the same as those in eqs. (4.31). For example, we can take $\varphi_i^{(n)}$ as

$$\varphi_i^{(n)}(k, l) = p_i^n (1 - p_i a)^{-k} \left(1 - \frac{1}{p_i} b\right)^{-l} e^{\eta_i} + q_i^n (1 - q_i a)^{-k} \left(1 - \frac{1}{q_i} b\right)^{-l} e^{\xi_i}, \quad (5.4)$$

$$\eta_i = \frac{1}{p_i} y + p_i x + \eta_{i0}, \quad (5.5a)$$

$$\xi_i = \frac{1}{q_i} y + q_i x + \xi_{i0}, \quad (5.5b)$$

where p_i , q_i and η_{i0} , ξ_{i0} are arbitrary constants which correspond to the wave numbers and phase parameters of solitons, respectively. It can easily be seen that $\varphi_i^{(n)}$ in eq. (5.4) actually satisfies eqs. (5.3).

The reduction is the technique to derive subhierarchies of equations by restricting the solution space, in other words, imposing some conditions to the solutions. In the case of RT equation, we take in eq. (5.4)

$$q_i = \frac{1}{p_i}, \quad (5.6)$$

and

$$b = a. \quad (5.7)$$

On the conditions (5.6) and (5.7), $\varphi_i^{(n)}$ in eq. (5.4) satisfies

$$(\partial_x + \partial_y) \varphi_i^{(n)} = \left(p_i + \frac{1}{p_i}\right) \varphi_i^{(n)}, \quad (5.8a)$$

$$\varphi_i^{(n)}(k-1, l-1) = (1 - p_i a) \left(1 - \frac{1}{p_i} a\right) \varphi_i^{(n)}(k, l). \quad (5.8b)$$

Hence we obtain

$$(\partial_x + \partial_y) \tau_n = S \tau_n, \quad (5.9a)$$

$$\tau_n(k-1, l-1) = P \tau_n(k, l), \quad (5.9b)$$

where

$$S = \sum_{i=1}^N (p_i + \frac{1}{p_i}), \quad (5.10a)$$

$$P = \prod_{i=1}^N (1 - p_i a)(1 - \frac{1}{p_i} a). \quad (5.10b)$$

By using eqs. (5.9), eqs. (5.1a) and (5.1c) are rewritten as

$$D_x^2 \tau_n(k, l) \cdot \tau_n(k, l) = 2(\tau_{n+1}(k, l)\tau_{n-1}(k, l) - \tau_n(k, l)\tau_n(k, l)), \quad (5.11a)$$

$$\begin{aligned} & \tau_n(k, l)\tau_n(k, l) - \tau_n(k+1, l)\tau_n(k-1, l) \\ &= a^2(\tau_n(k, l)\tau_n(k, l) - \tau_{n+1}(k+1, l)\tau_{n-1}(k-1, l)). \end{aligned} \quad (5.11b)$$

Here we may drop y and l -dependence. Thus eqs. (5.11) and (5.1b) give

$$D_x^2 \tau_n(k) \cdot \tau_n(k) = 2(\tau_{n+1}(k)\tau_{n-1}(k) - \tau_n(k)\tau_n(k)), \quad (5.12a)$$

$$(aD_x - 1)\tau_n(k) \cdot \tau_{n-1}(k-1) + \tau_n(k-1)\tau_{n-1}(k) = 0, \quad (5.12b)$$

$$\tau_n(k+1)\tau_n(k-1) - \tau_n(k)\tau_n(k) = a^2(\tau_{n+1}(k+1)\tau_{n-1}(k-1) - \tau_n(k)\tau_n(k)). \quad (5.12c)$$

Equations (5.12a) and (5.12c) are nothing but the bilinear form of the one-dimensional TL and DTL equations. From eqs. (5.2) and (5.4)-(5.7), the Casorati determinant solution of eqs. (5.12) is

$$\tau_n(k) = \begin{vmatrix} \varphi_1^{(n)}(k) & \varphi_1^{(n+1)}(k) & \cdots & \varphi_1^{(n+N-1)}(k) \\ \varphi_2^{(n)}(k) & \varphi_2^{(n+1)}(k) & \cdots & \varphi_2^{(n+N-1)}(k) \\ \vdots & \vdots & \ddots & \vdots \\ \varphi_N^{(n)}(k) & \varphi_N^{(n+1)}(k) & \cdots & \varphi_N^{(n+N-1)}(k) \end{vmatrix}, \quad (5.13)$$

$$\varphi_i^{(n)}(k) = p_i^n (1 - p_i a)^{-k} e^{\eta_i} + (\frac{1}{p_i})^n (1 - \frac{1}{p_i} a)^{-k} e^{\xi_i}, \quad (5.14)$$

$$\eta_i = p_i x + \eta_{i0}, \quad (5.15a)$$

$$\xi_i = \frac{1}{p_i} x + \xi_{i0}. \quad (5.15b)$$

Letting

$$f_n = \tau_n(n), \quad (5.16a)$$

$$g_n = \tau_{n-1}(n), \quad (5.16b)$$

$$\bar{g}_n = \tau_{n+1}(n), \quad (5.16c)$$

eqs. (5.12) with $k = n$ become the bilinear form of RT (2.1), and eq. (5.13) gives its solution (3.1). Thus we have completed the proof that the N -soliton solution for the RT equation is given by the Casorati determinant in eqs. (3.1). We note that besides the soliton solution, other solutions such as the rational exponential soliton can be obtained in the same way.

VI. CONCLUDING REMARKS

We have shown that the solution of the RT equation is given by the Casorati determinant. The composition of three Toda systems derives the relativistic modification of the TL equation. This fact means that the Toda equation has quite fruitful content.

The difference between the relativistic and non-relativistic TL's is only the choice of dependent variables in the D2DTL system. In the two-dimensional space of discrete independent variables n and k , we have the Casorati determinant solution $\tau_n(k)$. If we take $\tau_n(n)$ as the dependent variable f_n , we obtain the relativistic version of TL through the variable transformation (2.2a). The difference interval a gives the light speed c . On the other hand, adopting $\tau_n(k)$ with fixed k as the dependent variable, we get the original TL in the same variable transformation. Namely, the RT is a sublattice of the D2DTL while the TL is another sublattice.

It is well known that the other soliton equations such as the KdV, modified KdV, sine-Gordon and nonlinear Schrödinger equations are closely related to or derived from the Toda equation by suitable limiting procedures. Therefore we expect that the relativistic version of these equations can also be derived from the RT equation. We conjecture how to construct such relativistic soliton equations and their solutions.

- (1) For a given soliton equation, consider its integrable discretization.
- (2) Derive the proper two-dimensional lattice system (before reduction).
- (3) Take the different sublattice, that is, the different direction of propagation.
- (4) In a suitable variable transformation, the relativistic equation corresponding to the original one will be obtained.

It may be interesting to investigate the relativistic version of nonlinear integrable equations in this scenario.

It should be noted that the RT equation has many other aspects and there are several types of determinant solutions besides the Casorati one. Actually the Wronskian of two-directional type yields quite different solutions from the present N -soliton solution. We can

derive some interesting solutions from the Wronskian, which will be reported elsewhere.

ACKNOWLEDGMENT

The authors are grateful to Professor J. Hietarinta and Professor R. Hirota for valuable discussions.

- ¹ S. N. M. Ruijsenaars, Commun. Math. Phys. **133**, 217 (1990).
- ² M. Bruschi and O. Ragnisco, Phys. Lett. A **129**, 21 (1988); **134**, 365(1989).
- ³ Yu. B. Suris, Phys. Lett. A **145**, 113 (1990).
- ⁴ R. Hirota, *Solitons*, ed. by R. K. Bullough and P. J. Caudrey (Springer, Berlin, 1980).
- ⁵ J. Satsuma, J. Phys. Soc. Jpn. **46**, 359 (1979).
- ⁶ N. C. Freeman and J. J. C. Nimmo, Phys. Lett. A **95**, 1 (1983); J. J. C. Nimmo, Phys. Lett. A **99**, 281 (1983); N. C. Freeman, IMA J. Appl. Math. **32**, 125 (1984).
- ⁷ R. Hirota, M. Ito and F. Kako, Prog. Theor. Phys. Suppl. No. 94, 42 (1988).
- ⁸ M. Sato, RIMS Kokyuroku 439, 30 (1981).
- ⁹ M. Sato and Y. Sato, *Nonlinear Partial Differential Equations in Applied Science*, ed. H. Fujita, P. D. Lax and G. Strang (Kinokuniya/North-Holland, Tokyo, 1983), p.259.
- ¹⁰ E. Date, M. Kashiwara, M. Jimbo and T. Miwa, *Non-linear Integrable Systems — Classical Theory and Quantum Theory*, ed. M. Jimbo and T. Miwa (World Scientific, Singapore, 1983), p.39.
- ¹¹ M. Jimbo and T. Miwa, Publ. RIMS, Kyoto Univ. **19**, 943 (1983).
- ¹² T. Miwa, Proc. Jpn. Acad. **58A**, 9 (1982).
- ¹³ K. Ueno and K. Takasaki, *Group Representation and Systems of Differential Equations*, Adv. Stud. in Pure Math. **4** (Kinokuniya, Tokyo, 1984), p.1.
- ¹⁴ J. Hietarinta and J. Satsuma, Phys. Lett. A **161**, 267 (1991).
- ¹⁵ J. Matsukidaira, J. Satsuma and W. Strampp, Phys. Lett. A **147**, 467 (1990); J. Matsukidaira and J. Satsuma, J. Phys. Soc. Jpn. **59**, 3413 (1990); Phys. Lett. A **154**, 366 (1991); J. Hietarinta, K. Kajiwara, J. Matsukidaira and J. Satsuma, *Nonlinear Evolution Equations and Dynamical Systems*, ed. by M. Boiti, L. Martina and F. Penpinelli (World Scientific, Singapore, 1992).
- ¹⁶ Y. Ohta, R. Hirota, S. Tsujimoto and T. Imai, to be published in J. Phys. Soc. Jpn.

- ¹⁷ R. Hirota, Y. Ohta and J. Satsuma, Prog. Theor. Phys. Suppl. No. 94, 59 (1988).
- ¹⁸ R. Hirota and J. Satsuma, Prog. Theor. Phys. Suppl. No. 59, 64(1976).
- ¹⁹ R. Hirota, J. Phys. Soc. Jpn. **50**, 3785 (1981).